arXiv:1006.4316v1 [math.CA] 22 Jun 2010

JACOB'S LADDERS AND THE OSCILLATIONS OF THE FUNCTION $|\zeta(1/2+it)|^2$ AROUND ITS MEAN-VALUE; LAW OF THE ALMOST EXACT EQUALITY OF CORRESPONDING AREAS

JAN MOSER

Abstract. The oscillations of the function $Z^2(t),\ t\in[0,T]$ around the main part $\sigma(T)$ of its mean-value are studied in this paper. It is proved that an almost equality of the corresponding areas holds true. This result cannot be obtained by methods of Balasubramanian, Heath-Brown and Ivic.

1. Introduction

1.1. The Titchmarsh-Kober-Atkinson (TKA) formula

$$\int_0^\infty Z^2(t)e^{-2\delta t}\mathrm{d}t=\frac{c-\ln(4\pi\delta)}{2\sin\delta}+\sum_{n=0}^N c_n\delta^n+\mathcal{O}(\delta^{N+1}) \tag{1.1}$$

(see [17], p. 131) remained as an isolated result for the period of 56 years. We have discovered (see [5]) the nonlinear integral equation

$$\int_0^{\mu[x(T)]} Z^2(t)e^{-\frac{2}{x(T)}t}\mathrm{d}t=\int_0^T Z^2(t)\mathrm{d}t \tag{1.2}$$

in which the essence of the TKA formula is encoded. Namely, we have shown in [5] that the following almost exact expression for the Hardy-Littlewood integral

$$\int_0^T Z^2(t)\mathrm{d}t=\frac{\varphi(T)}{2}\ln\frac{\varphi(T)}{2}+(c-\ln(2\pi))\frac{\varphi(T)}{2}+c_0+\mathcal{O}\left(\frac{\ln T}{T}\right) \tag{1.3}$$

takes place, where $\varphi(T)$ is the Jacob's ladder, i.e. an arbitrary solution to the nonlinear integral equation (1.2).

Remark 1. Our formula (1.3) for the Hardy-Littlewood integral

$$\int_1^T\left|\zeta\left(\frac{1}{2}+it\right)\right|^2\mathrm{d}t=\int_1^T Z^2(t)\mathrm{d}t \tag{1.4}$$

has been obtained after the time period of 90 years since this integral appeared in 1918 with the first result

$$\int_1^T\left|\zeta\left(\frac{1}{2}+it\right)\right|^2\mathrm{d}t\sim T\ln T$$

(see [3], pp. 122, 151-156).

Key words and phrases. Riemann zeta-function.

1.2. Let us remind that

(A) The Good's Ω - theorem (see [2]) implies for the Balasubramanian formula (see [1])

$$\int_0^T Z^2(t)\mathrm{d}t \sim T\ln T + (2c-1-\ln 2\pi)T + R(T),\ R(T)=\mathcal{O}(T^{1/3+\epsilon}) \tag{1.5}$$

that

$$\limsup_{T\to\infty} |R(T)| = +\infty, \tag{1.6}$$

i.e. the error term in (1.5) is unbounded at $T\to\infty$.

(B) In the case of our formula (1.3) the error term definitely tends to zero

$$\lim_{T\to\infty} r(T)=0; \qquad r(T)=\mathcal{O}\left(\frac{\ln T}{T}\right), \tag{1.7}$$

i.e. our formula is almost exact (see [5]).

Remark 2. In this paper the geometric interpretation of (1.6) and (1.7) is obtained.

1.3. For the mean-value of the function $\left|\zeta\left(\frac{1}{2}+it\right)\right|^2 = Z^2(t)$, where

$$Z(t)=e^{i\vartheta(t)}\zeta\left(\frac{1}{2}+it\right), \quad \vartheta(t)=-\frac{t}{2}\ln\pi + \mathrm{Im}\ln\Gamma\left(\frac{1}{4}+i\frac{t}{2}\right),$$

we obtain from (1.3)

$$\frac{1}{T}\int_0^T Z^2(t)\mathrm{d}t = \frac{\varphi(T)}{2T}\ln\frac{\varphi(T)}{2} + (c-\ln 2\pi)\frac{\varphi(T)}{2T} + \frac{c_0}{T} + \mathcal{O}\left(\frac{\ln T}{T}\right). \tag{1.8}$$

Let

$$\sigma(T) = \frac{\varphi(T)}{2T}\ln\frac{\varphi(T)}{2} + (c-\ln 2\pi)\frac{\varphi(T)}{2T} + \frac{c_0}{T} \tag{1.9}$$

denote the main part of the mean-value (1.8). In this paper the oscillation of the values of the function $Z^2(t),\ t\in[0,T]$ around the main part $\sigma(T)$ of its mean-value are studied.

Remark 3. The main result of this paper is the following statement: the areas of the figures corresponding to the parts of the graph of the function $Z^2(t),\ t\in[0,T]$ given by inequalities $Z^2(t)\geq\sigma(T)$ and $Z^2(t)\leq\sigma(T)$, respectively, are almost exactly equal.

This paper is a continuation of the series [5]-[16].

2. RESULT

2.1. Let (see (1.9))

$$\begin{aligned} S^+(T) &= \{t:\ Z^2(t)\geq\sigma(T),\ t\in[0,T]\}, \\ S^-(T) &= \{t:\ Z^2(t)<\sigma(T),\ t\in[0,T]\} \end{aligned} \tag{2.1}$$

and

$$\begin{aligned} \Pi^+(T) &= \{(t,y):\ \sigma(T)\leq y\leq Z^2(t),\ t\in S^+(T)\}, \\ \Pi^-(T) &= \{(t,y):\ Z^2(T)\leq y\leq\sigma(t),\ t\in S^-(T)\}, \end{aligned} \tag{2.2}$$

i.e. Π^+ is the figure that corresponds to the parts of the graph of $y=Z^2(t),\ t\in[0,T]$ lying above the segment $y=\sigma(T)$ and similarly Π^- corresponds to the parts of the graph lying under that segment. Let $m\{\Pi^+(T)\},\ m\{\Pi^-(T)\}$ denote measures of corresponding figures, i.e.

$$\begin{aligned} m\{\Pi^+(T)\} &= \int_{S^+(T)}\{Z^2(t)-\sigma(T)\}\mathrm{d}t, \\ m\{\Pi^-(T)\} &= \int_{S^-(T)}\{\sigma(T)-Z^2(t)\}\mathrm{d}t. \end{aligned} \tag{2.3}$$

The following theorem holds true.

Theorem. First of all, we have the formula

$$m\{\Pi^+(T)\}=m\{\Pi^-(T)\}+\mathcal{O}\left(\frac{\ln T}{T}\right) \tag{2.4}$$

(see (1.3), (1.9), (2.1)-(2.3)). Next, the structure of the formula (2.4) is as follows: there are the functions $\eta_1(T),\ \eta_2(T)$ that the following formulae

$$\begin{aligned} m\{\Pi^+(T)\} &= \frac{1+o(1)}{2\pi^2}\frac{T\ln^4 T}{\eta_1-\eta_2}-\frac{\eta_2}{\eta_1-\eta_2}\mathcal{O}\left(\frac{\ln T}{T}\right), \\ m\{\Pi^-(T)\} &= \frac{1+o(1)}{2\pi^2}\frac{T\ln^4 T}{\eta_1-\eta_2}-\frac{\eta_1}{\eta_1-\eta_2}\mathcal{O}\left(\frac{\ln T}{T}\right) \end{aligned} \tag{2.5}$$

hold true, and

$$AT^{2/3}\ln^4 T< m\{\Pi^+(T)\}, m\{\Pi^-(T)\}<AT\ln T. \tag{2.6}$$

In addition to (2.6): on the Lindelöf hypothesis

$$m\{\Pi^+(T)\}, m\{\Pi^-(T)\}>A(\epsilon)T^{1-\epsilon}, \tag{2.7}$$

and on Riemann hypothesis

$$m\{\Pi^+(T)\}, m\{\Pi^-(T)\}>T^{1-\frac{A}{\ln\ln T}}. \tag{2.8}$$

Corollary. We have by (2.5), (2.6)

$$\eta_1(T)-\eta_2(T)>A\ln^3 T. \tag{2.9}$$

Remark 4. Since from (2.4)

$$\lim_{T\to\infty}[m\{\Pi^+(T)\}-m\{\Pi^-(T)\}]=0 \tag{2.10}$$

follows then we have the almost exact equality of the areas $m\{\Pi^+(T)\}$ and $m\{\Pi^-(T)\}$.

2.2. In the case of the Balasubramanian formula (1.5) we have (comp. (1.3), (1.9))

$$\sigma_1(T)=\ln T+2c-1-\ln 2\pi.$$

Let $S_1^+(T), S_1^-(T), \Pi_1^+(T), \Pi_1^-(T), m\{\Pi_1^+(T)\}, m\{\Pi_1^-(T)\}$ correspond to $\sigma_1(T)$ similarly to (2.1)-(2.3). Then from (1.5) we obtain

$$m\{\Pi_1^+(T)\}=m\{\Pi_1^-(T)\}+\mathcal{O}(T^{1/3+\epsilon}),\ T\to\infty, \tag{2.11}$$

and (see (1.6)

$$\limsup_{T\to\infty}|m\{\Pi_1^+(T)\}-m\{\Pi_1^-(T)\}|=+\infty. \tag{2.12}$$

Remark 5. The following holds true:

(A) Our formula (1.3) which has been obtained by means of the Jacob's ladders leads to the almost exact equality of the areas (see (2.4), (2.10).

(B) The Balasubramanian formula (1.5) which has been obtained by means of estimation of trigonometric sums leads to the formula (2.11) that possesses quite large uncertainty (2.11) and this error term cannot be removed.

3. PROOF OF THEOREM

3.1. We obtain from (1.3), (2.1)

$$\int_{S^+(T)}\{Z^2(t)-\sigma(T)\}\mathrm{d}t+\int_{S^-(T)}\{Z^2(t)-\sigma(T)\}\mathrm{d}t=\mathcal{O}\left(\frac{\ln T}{T}\right) \tag{3.1}$$

and from (3.1) by (2.3) the formula

$$m\{\Pi^+(T)\}-m\{\Pi^-(T)\}=\mathcal{O}\left(\frac{\ln T}{T}\right) \tag{3.2}$$

follows, i.e. (2.4).

3.2. Next, from the Ingham formula (see [4], p. 277, [17], p. 125)

$$\int_0^T Z^4(t)\mathrm{d}t=\frac{1}{2\pi^2}T\ln^4 T+\mathcal{O}(T\ln^3 T) \tag{3.3}$$

we obtain (see (1.9)

$$\int_0^T\{Z^4(t)-\sigma^2(T)\}\mathrm{d}t=\frac{1}{2\pi^2}T\ln^4 T-T\sigma^2(T)+\mathcal{O}(T\ln^3 T). \tag{3.4}$$

Since ($\varphi(T)\sim T$)

$$T\sigma^2(T)=\mathcal{O}\left\{\frac{\varphi^2(T)}{T}\ln^2\frac{\varphi(T)}{2}\right\}=\mathcal{O}(T\ln^2 T),$$

then from (3.4) the formula

$$\int_0^T\{Z^4(t)-\sigma^2(T)\}\mathrm{d}t=\frac{1+o(1)}{2\pi^2}T\ln^4 T \tag{3.5}$$

follows.

3.3. Since $Z^4(t)-\sigma^2(T)=(Z^2-\sigma)(Z^2+\sigma)$ and $Z^2(t)-\sigma(T)$ is always of the same sign on $S^+(T)$ and on $S^-(T)$, respectively, then from (3.5) we obtain (see (2.3))

$$\eta_1(T)m\{\Pi^+(T)\}-\eta_2(T)m\{\Pi^-(T)\}=\frac{1+o(1)}{2\pi^2}T\ln^4 T, \tag{3.6}$$

where $\eta_1=\eta_1(T),\ \eta_2=\eta_2(T)$ are the mean-values of $Z^2(t)+\sigma(T)$ relatively to the values of the functions $Z^2(t)-\sigma(T)$ and $\sigma(T)-Z^2(t)$, respectively on the sets $S^+(T)$ and $S^-(T)$, respectively. It is clear that

$$A\ln T<\eta_1(T),\eta_2(T)<AT^{1/3} \tag{3.7}$$

(see (1.9); $|Z(t)|<t^{1/6}$). Next, $\eta_1(T)\neq\eta_2(T)$ is also true. Since if $\eta_1=\eta_2$ then by (3.2), (3.6), (3.7) we would have the contradiction. Hence, from the simple system of linear equations (3.2), (3.6) we obtain (2.5).

3.4. Since

$$0 < \eta_1 - \eta_2 < \eta_1 < AT^{1/3} \tag{3.8}$$

(see (2.5), (3.7)) then we obtain from (2.5) the lower estimates in (2.6). Next we have (see (1.9), (2.3)

$$m\{\Pi^+(T)\}, m\{\Pi^-(T)\} < \int_0^T \{Z^2(t) - \sigma(T)\}\mathrm{d}t < AT\ln T$$

i.e. the upper estimates in (2.6) hold true.

3.5. Following the Lindelöf and the Riemann conjectures the estimates

$$Z^2(t) < A(\epsilon)t^\epsilon, \quad t^{\frac{A}{\ln\ln t}}$$

take place correspondingly and then the conditional estimates (2.7) and (2.8) follow.

I would like to thank Michal Demetrian for helping me with the electronic version of this work.

References

[1] R. Balasubramanian, ‘An improvement on a theorem of Titchmarsh on the mean square of $|\zeta(1/2+it)|^2$‘, Proc. London Math. Soc. 3, 36 (1978), 540-575.

[2] A. Good, ‘Ein Ω - resultat für quadratische Mittel der Riemannschen Zetafunktion auf der kritische Linie‘, Invent. Math. 41, (1977), 233-251.

[3] G.H. Hardy and J.E. Littlewood, ‘Contribution to the theory of the Riemann zeta-function and the theory of the distribution od Primes‘, Acta Math. 41, (1918), 119-195.

[4] A.E. Ingham, ‘Mean-value theorems in the theory of the Riemann zeta-function‘, Proc. Lond. Math. Soc. (2), 27, (1926), 273-300.

[5] J. Moser, ‘Jacob's ladders and the almost exact asymptotic representation of the Hardy-Littlewood integral', (2008), arXiv:0901.3973.

[6] J. Moser, ‘Jacob's ladders and the tangent law for short parts of the Hardy-Littlewood integral', (2009), arXiv:0906.0659.

[7] J. Moser, ‘Jacob's ladders and the multiplicative asymptotic formula for short and microscopic parts of the Hardy-Littlewood integral', (2009), arXiv:0907.0301.

[8] J. Moser, ‘Jacob's ladders and the quantization of the Hardy-Littlewood integral', (2009), arXiv:0909.3928.

[9] J. Moser, ‘Jacob's ladders and the first asymptotic formula for the expression of the sixth order $|\zeta(1/2+i\varphi(t)/2)|^4|\zeta(1/2+it)|^2$', (2009), arXiv:0911.1246.

[10] J. Moser, ‘Jacob's ladders and the first asymptotic formula for the expression of the fifth order $Z[\varphi(t)/2+\rho_1]Z[\varphi(t)/2+\rho_2]Z[\varphi(t)/2+\rho_3]\hat{Z}^2(t)$ for the collection of disconnected sets‘, (2009), arXiv:0912.0130.

[11] J. Moser, ‘Jacob's ladders, the iterations of Jacob's ladder $\varphi_1^k(t)$ and asymptotic formulae for the integrals of the products $Z^2[\varphi_1^n(t)]Z^2[\varphi^{n-1}(t)]\cdots Z^2[\varphi_1^0(t)]$ for arbitrary fixed $n \in \mathbb{N}$‘ (2010), arXiv:1001.1632.

[12] J. Moser, ‘Jacob's ladders and the asymptotic formula for the integral of the eight order expression $|\zeta(1/2+i\varphi_2(t))|^4|\zeta(1/2+it)|^4$‘, (2010), arXiv:1001.2114.

[13] J. Moser, ‘Jacob's ladders and the asymptotically approximate solutions of a nonlinear diophantine equation‘, (2010), arXiv: 1001.3019.

[14] J. Moser, ‘Jacob's ladders and the asymptotic formula for short and microscopic parts of the Hardy-Littlewood integral of the function $|\zeta(1/2+it)|^4$‘, (2010), arXiv:1001.4007.

[15] J. Moser, ‘Jacob's ladders and the nonlocal interaction of the function $|\zeta(1/2+it)|$ with $\arg\zeta(1/2+it)$ on the distance $\sim (1-c)\pi(t)$‘, (2010), arXiv: 1004.0169.

[16] J. Moser, ‘Jacob's ladders and the $\tilde{Z}^2$ - transformation of polynomials in $\ln\varphi_1(t)$‘, (2010), arXiv: 1005.2052.

[17] E.C. Titchmarsh, ‘The theory of the Riemann zeta-function‘, Clarendon Press, Oxford, 1951.

Department of Mathematical Analysis and Numerical Mathematics, Comenius University, Mlynska Dolina M105, 842 48 Bratislava, SLOVAKIA
E-mail address: `jan.mozer@fmph.uniba.sk`